



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

REMARK. In (I) the sign changes with either x or y , that is, at the ends of either axis; while in (II) the sign changes only at the ends of the major axis. The factor $ex+a$ vanishes only on the left-hand directrix; hence not in the ellipse.

Also solved by G. B. M. Zerr, and J. Scheffer.

246. Proposed by A. M. HARDING, Adjunct Professor, University of Arkansas, Fayetteville, Ark.

A pentagon $ABCDE$, formed of equal uniform heavy rods connected by smooth joints at their ends, is supported symmetrically in a vertical plane with A uppermost, and AB and AE in contact with two smooth pegs in the same horizontal line. Prove that if the pentagon is regular, the pegs must divide AB and AE each in the ratio $1+\sin(\pi/10):3\sin(\pi/10)$. *Jeans' Theoretical Mechanics*, page 112, number 13.

Solution by PROF. F. L. GRIFFIN, Ph. D., Williams College.

Denote by P the point of contact of AB with one peg, and let $AP=2m$ and $PB=2n$. Also let the weight of each rod be $2W$, and the resistance exerted by each peg be R .

Consider first the rod AB , making an angle of 36° with the horizontal. At A the reaction of AE is horizontal (say X_a); for if there were a vertical component, then AB would exert upon AE an oppositely directed vertical component, contrary to the hypothesis of symmetry. At B the reaction of BC is unknown; denote its components by X_b and Y_b . The other forces applied to AB are its weight $2W$, and the reaction R inclined 54° to the horizontal. Equating to zero the sums of the vertical components, horizontal components, and moments about P we obtain (since the distance from P to the midpoint of AB is $2m-(m+n)=m-n$).

$$\begin{aligned} (1), (2) \quad & Y_b + R\sin 54^\circ - 2W = 0, \quad X_a + X_b + R\cos 54^\circ = 0, \\ (3) \quad & X_b 2n\cos 54^\circ + Y_b 2n\sin 54^\circ + 2W(m-n)\sin 54^\circ - X_a 2m\cos 54^\circ = 0. \end{aligned}$$

Now R is easily found by considering the pentagon as a whole, to which are applied three external forces, R , R and the weight $10W$. The vanishing of the vertical component gives

$$(4) \quad 2R\sin 54^\circ - 10W = 0, \quad R = 5W\csc 54^\circ.$$

To obtain X_b consider the rod BC , to which is applied at B the reaction of AB whose components are respectively $(-X_b)$ and $(-Y_b)$. Equating to zero the resultant moment about C , we have, since $BC=2(m+n)$ and is inclined 72° to the horizontal:

$$(5) \quad (-Y_b) \cdot 2(m+n)\sin 18^\circ - (-X_b) 2(m+n)\cos 18^\circ - 2W(m+n)\sin 18^\circ = 0.$$

Using (4) to solve successively (1), (5), and (2) we obtain

$$Y_b = -3W, \quad X_b = (W + Y_b)\tan 18^\circ = -2W\tan 18^\circ, \\ X_a = 2W\tan 18^\circ - 5W\cot 54^\circ.$$

Putting these values into (3) and dividing by $2W$ we get

$$(6) \quad -2\tan 18^\circ \cdot n \cdot \cos 54^\circ - 3W \cdot n \sin 54^\circ + (m - n) \sin 54^\circ \\ + (5\cot 54^\circ - 2\tan 18^\circ) \cdot m \cos 54^\circ = 0.$$

Multiplying (6) by $\sin 54^\circ$, collecting m and n , and later employing the identities $2\sin^2 x = 1 - \cos 2x$, $2\cos^2 x = 1 + \cos 2x$, and $2\sin x \cos x = \sin 2x$, we obtain:

$$m[\sin^2 54^\circ - 2\sin 54^\circ \cos 54^\circ \tan 18^\circ + 5\cos^2 54^\circ] \\ = n[4\sin^2 54^\circ + 2\sin 54^\circ \cos 54^\circ \tan 18^\circ], \\ m[3 + 2\cos 108^\circ - \sin 108^\circ \tan 18^\circ] = n[2 - 2\cos 108^\circ + \sin 108^\circ \tan 18^\circ], \\ \text{or, } m[3 - 3\sin 18^\circ] = n[2 + 3\sin 18^\circ],$$

$$\text{whence, } m : n = 2 + 3\sin(\pi/10) : 3 - 3\sin(\pi/10).$$

The proposed result follows immediately from this one, if we recall from geometry that the half-side of the regular inscribed decagon $= R(\sqrt{5} - 1)/4$, so that $\sin 18^\circ$ satisfies the equation

$$(4\sin x + 1)^2 = 5 \text{ or } 4\sin^2 x + 2\sin x = 1.$$

For from this we obtain $\sin x(3\sin x + 2) = 1 - \sin^2 x$, or $2 + 3\sin x : 3(1 - \sin x) :: 1 + \sin x : 3\sin x$.

PROBLEMS FOR SOLUTION.

ALGEBRA.

343. Proposed by THEODORE L. DeLAND, Treasury Department, Washington, D. C.

A, on contracting to execute a piece of work for \$300 and finding after working alone one day that he had finished but 1% of the entire work, engaged B to assist him at the beginning of the second day, with the understanding, that B on each day was to do 6% as much work as had been completed previously, while A each day was to do an amount of